LAWNPORN HISTORY

Product Information Sheet



What is Lawnporn History?

Lawnporn History Systemic insecticide is a professional turf solution for the control of African Black Beetle, Argentinean Scarabs and Bilbugs Larvae. The active ingredient, Thiamethoxam works through systemic action.



Features of Lawnporn History

- ✓ Control of African Black Beetle, Argentinian Scarab and Billbug larvae (grubs)
- ✓ Low to no odour
- Suitable for turf situations with irregular irrigation
- ✓ Mode of Action is Group 4A Neonicotinoids
- ✓ Ideal for the home lawn enthusiast
- Refer to application chart for directions

Active Ingredients

Thiamethoxam 250g/L

> Insect Management

DIRECTIONS FOR USE

SITUATION	PESTS	RATE	CRITICAL COMMENTS
Turf, Lawns	1st instar larvae of African Black Beetle (Heteronychus arator)	800mL to 1.2L / ha in at least 500L water per ha or 8mL to 12mL / 100m² In at least 5L water per 100m²	Apply prior to or at peak egg hatch, mid spring to early summer, or when small larvae are present. Ensure product placement as close to soil surface as possible. Total application volume should not be lower than 500L / ha. For best results use extremely coarse droplets [Turbo Floodjet* (TF5) or TurfJet* (TTJ10)]. Preferably spray onto wet or dewy grass. Irrigate with 6 to 12 mm of water commencing within 1 hour of application.
	Larvae of Billbug (Sphenophorus brunnipennis)		Monitor adult activity through late spring and early summer. Spray when numbers peak or when small larvae (4 mm) are found in the thatch or surface soil. Early application is essential to minimise grass damage due to feeding. Use the higher rate when pest pressure is high or 2nd instar larvae are present. Ensure product placement as close to soil surface as possible. Total application volume should not be lower than 500L / ha. For best results use extremely coarse droplets [Turbo Floodjet* (TF5) or TurfJet* (TTJ10)]. Preferably spray onto wet or dewy grass. Irrigate with 6 to 12 mm of water commencing within 1 hour of application.

WITHHOLDING PERIODS

DO NOT GRAZE TREATED TURF/LAWN OR FEED TURF/LAWN
CLIPPINGS FROM ANY TREATED AREA TO POULTRY OR

APPLICATION

To be effective Lawnporn History requires to be placed as close to the soil surface as possible. Ensure that equipment is properly calibrated to give an even distribution at the correct volume. Apply in a minimum of 500L water/ha. For best results use extremely coarse droplets [Turbo Floodjet* (TF5) or TurfJet* (TTJ10)].

> First Aid & Safety Directions

In case of contact with eyes or skin, flush at once with large amounts of water. If irritation persists, seek medical attention. If swallowed, drink large amountsof water and induce vomiting. Seek medical attention.

FOR MORE INFORMATION SPEAK TO YOUR AMGROW TERRITORY MANAGER.

