

PROPYZAMIDE 500

HERBICIDE

500 g/L PROPYZAMIDE

for **Winter grasses**



patersons curse



control

SureFire

PROPYZAMIDE 500

HERBICIDE

500 g/L PROPYZAMIDE

SureFire PROPYZAMIDE 500 herbicide is a member of the benzamides, a Group D herbicide. It is selective, systemic and is absorbed by roots and translocated throughout the plant.

It is a residual, soil acting herbicide, suitable as either a pre-emergent or post-emergent means of control of winter grass and many broadleaf weeds (see table below).

Best results are obtained when applied during cooler weather. Rain or irrigation is necessary to ensure optimal results with this product.

Not recommended to mix product with any other pesticides.

SITUATION	WEEDS CONTROLLED	STATE	RATE L/ha	CRITICAL COMMENTS
Lettuce Direct sown	Barnyard Grass, Blackberry, Nightshade, Chickweed, Portulaca, Winter Grass, Ryegrass, Summer Grass, Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles	All States	4.5 L in 330 – 550 L water	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.
		Vic Only	2.3 L in 300-550 L water	Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.
Turfs & Lawns	Winter grass			Rate Bowling Green 37 x 37 m
Common Couch		All States	1.2 L (12 mL/100 m ²)	170 mL
Queensland Blue couch		Qld only	0.6 L (6 mL/100 m ²)	85 mL
Buffalo grass		WA only	1.2 L (12 mL/100 m ²)	170 mL
				Apply during autumn and Winter, if established turf is infested with seedling winter grass. Ensure a uniform coverage. Irrigate immediately after application with 12-25 mm irrigation or rain and again 24 hours later. Repeat if necessary on Qld. Blue Couch after 4 weeks. DO NOT re-seed for 60 days after treatment. Use ONLY on established turf. DO NOT use on Bluegrass, ryegrass, fescue or bent grass lawns.
Lucerne, Clover, Medics etc grown for forage, hay or seed production	Annual Ryegrass, Winter grass, Barley grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue and Rats Tail Fescue	NSW, ACT, SA, WA, Tas only	1-1.5 L	SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July WEEDS: Apply when weeds are at post or pre-emergent stages.
			1.5-2 L	BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25 mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatments is recommended.

Container sizes: Carton quantity:

5 litre 4

APVMA Approval No: 70362 / 63278

GROUP D HERBICIDE

WITHHOLDING PERIODS: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.



© PCT International Pty Ltd

www.pctrural.com.au