Initiator Tablet	Page: 1 of 5
Approved text label – 2 nd March 2009	
Text above this line is not included in the label	

Front panel of label

CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Initiator®

Systemic Plant Insecticide and Fertiliser

ACTIVE CONSTITUENT: 200 g/kg IMIDACLOPRID

GROUP 4A INSECTICIDE

* Kg NET (* x 2.5 g

Tablets)

For use in the protection against insect damage and to provide enhanced growth in eucalyptus trees, roses, azaleas, lillypillies, palms and magnolias.

Guaranteed Minimum Fertiliser Analysis:

Nitrogen (N) (as slow release)	3.85%
Nitrogen (N) as Ammonium Nitrate	1.75%
Total Nitrogen (N):	5.6%
Phosphorous (P) (as calcium phosphate)	2.6%
Phosphorous (P) (Slow release)	2.8%
Total Phosphorous:	5.4%
Potassium (K) (Slow release):	4.2%
Magnesium (Mg) (Slow release):	1.4%

*APVMA Approval Nos.: 60391/3kg/0209 (1200 x 2.5 g tablets) 60391/625g-1.25kg/0209 (250-500 x 2.5 g tablets)

Bayer CropScience Pty Ltd A.B.N. 87 000 226 022 391-393 Tooronga Rd, Hawthorn East, Vic 3123 Telephone: 1800 804 479

DOM:

B/N:

Initiator Tablet	Page:	2 of 5
Approved text label – 2 nd March 2009		
Text above this line is not included in the label		

Front panel of label

CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Initiator®

Systemic Plant Insecticide and Fertiliser

ACTIVE CONSTITUENT: 200 g/kg IMIDACLOPRID

GROUP 4A INSECTICIDE

750g NET (300 x 2.5 g

Tablets)

For use in the protection against insect damage and to provide enhanced growth in eucalyptus trees, roses, azaleas, lillypillies, palms and magnolias.

Guaranteed Minimum Fertiliser Analysis:

85%
75%
6%
6%
8%
4%
2%
4%

APVMA Approval No.: 60391/625 g - 1.25 kg/0209

Bayer CropScience Pty Ltd A.B.N. 87 000 226 022 391-393 Tooronga Rd, Hawthorn East, Vic 3123 Telephone: 1800 804 479

DOM:

B/N:

Text above this line is not included in the label

Auxillary panel

r use		
PEST	RATE	CRITICAL COMMENTS
ised, newly planted and established plar protection and fertiliser effects the tablet newly established seedlings, containeriz	nts. has been sh	own to improve survival and
amental plants.		
Aphids Azalea lace bug	1 tablet per m of plant height.	Transplanted stock: Place tablet(s) in planting hole, immediately beneath root ball, at time of planting
Psyllids		Established in-ground or
Scale		potted plants: Create a hole in the soil approximately 5 cm deep, as close to the absorptive roots as possible (at least 10 cm away from stem of plant). Place tablet in hole and ensure tablet is completely covered with soil after placement. In plants greater than 1 m in height, where multiple tablets are required, space out the tablets evenly around the plant.
Various defoliating insect pests including Chrysomelid larvae and adults (eg. <i>Chrysophtharta</i> spp., <i>Paropsis</i> spp., <i>Cadmus</i> spp.), scarab pests (eg. <i>Heteronyx</i> <i>dimidiata</i> and <i>H. crinitus</i>) Suppression of damage caused by Autumn gum moth (<i>Mnesampela</i> <i>privata</i>) and Light brown apple moth (<i>Epiphyas postvittana</i>) Stem damaging scarab pests eg. adult African Black Beetle (<i>Heteronychus arator</i>) Sap-sucking pests eg. Psyllids and Gum-tree scale Root-feeding scarab pests eg.	1 tablet per seedling tree	Transplanted seedling stock: Place tablet in planting hole, immediately beneath or adjacent to root ball, at time of planting
	PEST will control or significantly reduce damagised, newly planted and established plar protection and fertiliser effects the tablet newly established seedlings, containerizamental plants. Aphids Azalea lace bug Psyllids Mealybugs Scale Various defoliating insect pests including Chrysomelid larvae and adults (eg. Chrysophtharta spp., Paropsis spp., Cadmus spp.), scarab pests (eg. Heteronyx dimidiata and H. crinitus) Suppression of damage caused by Autumn gum moth (Mnesampela privata) and Light brown apple moth (Epiphyas postvittana) Stem damaging scarab pests eg. adult African Black Beetle (Heteronychus arator)	PESTRATEwill control or significantly reduce damage caused by ised, newly planted and established plants. protection and fertiliser effects the tablet has been sh newly established seedlings, containerized trees and amental plants.Aphids1 tablet per m of plant height.Azalea lace bug1 tablet per m of plant height.Psyllids1 tablet per m of plant height.Various defoliating insect pests including Chrysomelid larvae and adults (eg. Chrysophtharta spp., Paropsis spp., Cadmus spp.), scarab pests (eg. Heteronyx dimidiata and H. crinitus)1 tablet per seedling treeSuppression of damage caused by Autumn gum moth (Mnesampela privata) and Light brown apple moth (Epiphyas postvittana)1 terested persectionStem damaging scarab pests eg. adult African Black Beetle (Heteronychus arator)Sap-sucking pests eg. Psyllids and Gum-tree scaleRoot-feeding scarab pests eg.Root-feeding scarab pests eg.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

<u>Auxillary panel – outer box</u>

GENERAL INSTRUCTIONS:

Initiator tablets are formulated to release active ingredient slowly. Release of active ingredient is dependent on presence of adequate soil moisture. For best results, irrigate the plant after treatment, and ensure that the plant receives regular irrigation for at least 2-3 months after treatment.

FORESTRY APPLICATIONS

Once the seedling has been placed into the planting hole where an Initiator tablet has been applied there is no requirement for any deviation from normal planting practice.

This tablet formulation has been shown to significantly improve growth and establishment rates of eucalypt plantations in areas where insect pests are significant limitations to growth and establishment.

PRODUCTION AND LANDSCAPE ORNAMENTALS

Initiator tablets can be used as directed on the plant species described above in areas such as production and retail nurseries, home gardens and residential complexes; business, office and shopping complexes; golf courses, airports, cemeteries, parks, playgrounds, athletic fields, and other landscaped recreational areas.

INSECTICIDE RESISTANCE WARNING: GROUP 4A INSECTICIDE

For insecticide resistance management Initiator is a Group 4A insecticide.

Some naturally occurring insect biotypes resistant to Initiator and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Initiator or other Group 4A insecticides are used repeatedly. The effectiveness of Initiator on resistant individuals could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience accepts no liability for any losses that may result from the failure of this product to control resistant insects. Initiator Tablets may be subject to specific resistance management strategies. For further information contact your local supplier or Bayer Environmental Science representative.

PRECAUTIONS

DO NOT allow stock to graze treated trees or pasture adjacent to trees.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT leave tablets exposed above ground as they are dangerous to wildlife, particularly birds.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight.

Triple rinse container before disposal. Dispose of rinsings in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. If recycling, replace lid and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. DO NOT burn empty containers or product. DO NOT re-use empty containers for any other purpose.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. Wear rubber gloves. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (131126).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd

Initiator Tablet Approved text label – 2 nd March 2009	Page:	
Text above this line is not included in the label		-

accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval Number: 60391/625g – 1.25 kg/0209

Initiator® is a Registered Trademark of the Bayer Group

Bayer CropScience Pty Ltd ABN 87 000 226 022 391-393 Tooronga Rd Hawthorn East, Vic. 3123 Telephone: 1800 804 479 www.environmentalscience.bayer.com.au