

## DIRECTIONS FOR USE

**RESTRAINTS: DO NOT TREAT** animals with this product more than 3 times in any given 12 month period. Allow at least 21 days between treatments. More frequent use of this product is likely to result in residues above the MRL.

**Precaution:** The safety of this product on pregnant cows or in calves has not been established.

ANIMAL AND TARGET PEST	DOSAGE	ADMINISTRATION
Cattle, Ready-to-Use Pour-on Application:	1mL / 30 kg body weight per animal up to 15mL maximum. Reapply as needed but not more often than once every 21 days	<b>SHAKE WELL BEFORE USE</b> Apply with a suitable applicator gun
Buffalo Flies ( <i>Haematobia imitans exigua</i> )		For Buffalo Fly control: Apply along the backline, from the shoulders to the tail butt, when fly numbers exceed acceptable levels
Cattle Lice, biting and sucking species ( <i>Bovicola bovis</i> , <i>Linognathus vituli</i> , <i>Haematopinus eurysternus</i> )		For Cattle Lice control: Apply along the backline, from the head poll down the neck and over the shoulders. For optimum residual control, reapply after 3 weeks.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY  
TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

## **DIRECTIONS FOR USE:**

### **SHAKE WELL BEFORE USE:**

Measure the required dose into the chamber and apply as a pour-on. Apply 1 mL per 30 kg bodyweight up to a maximum of 15 mL per animal. The maximum number of applications is 3 per season.

**FOR BUFFALO FLY CONTROL:** Apply along the backline, from the shoulders to the tail butt, when fly numbers exceed acceptable levels.

**FOR CATTLE LICE CONTROL:** Apply along the backline, from the head poll down the neck and over the shoulders. For optimum residual control, reapply after 3 weeks.

Reapply as needed but not more often than once every 21 days.

### **RESTRAINTS:**

**MEAT: DO NOT USE** less than 7 days before slaughter for human consumption.

**MILK: DO NOT USE** on lactating or pregnant cows where milk products may be used for human consumption.

**DO NOT TREAT** animals with this product more than 3 times in any given 12 month period. Allow at least 21 days between treatments. More frequent use of this product is likely to result in residues above the MRL.

**PRECAUTION:** The safety of this product on pregnant cows or in calves has not been established.

**DOSAGE AND ADMINISTRATION:** SHAKE WELL BEFORE USING: For the 1 Litre pack size, loosen cap and squeeze bottle to fill measuring chamber with required dosage. Remove cap and pour along animal's back-line.

1 mL/30 kg bodyweight per animal up to 15 mL maximum.

Reapply as needed but not more often once every 21 days. DO NOT apply more than 3 times in any 12 month period.

**GENERAL DIRECTIONS:** SWAT Pour-On for Cattle will give effective control of pyrethroid-susceptible Buffalo fly and Cattle Lice when used as directed. The level and duration of control may be reduced due to high fly pressure. Treat all animals in the group. The length of residual protection against Buffalo flies will vary depending on the level of insecticide resistance within a particular area. SWAT Pour-On is effective against biting lice and two species of sucking lice.

DO NOT TREAT animals with this product more than 3 times in any given 12 month period. Allow at least 21 days between treatments. More frequent use of this product is likely to result in residues above the MRL.

**WITHHOLDING PERIODS:**

**MEAT: DO NOT USE** less than 7 days before slaughter for human consumption.

**MILK: DO NOT USE** on lactating or pregnant cows where milk products may be used for human consumption.

**EXPORT OF TREATED PRODUCE: DO NOT USE** less than 14 days before slaughter for export. Before using this product, confirm the current ESI from Pharmachem on 07 3868 0333 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues)).

**SAFETY DIRECTIONS:** May irritate the eyes. Avoid contact with eyes. Wash hands after use.

**FIRST AID:** If poisoning occurs, contact a doctor or the Poisons Information Centre; Phone 131126.

**SAFETY DATA SHEET:** If additional hazard information is required, refer to the Safety Data Sheet which can be obtained by contacting Pharmachem.

**ENVIRONMENTAL STATEMENTS:** This product is highly toxic to fish. DO NOT contaminate streams, rivers or waterways with the insecticide or used containers.

**DISPOSAL:** Triple rinse container before disposal. Dispose of rinsate in a disposal pit specifically marked and set up for this purposes clear of waterways, vegetation and roots. DO NOT dispose of undiluted chemicals on site. Break, crush or puncture and bury container in a local authority landfill. If not available, bury the containers below 500mm in the disposal pit referred to above. Empty containers and product should not be burned.

**STORAGE:** Store below 30°C (Room temperature). Store upright in the original container tightly closed, in a dry place. DO NOT store in direct sunlight.

**WARRANTY:** The use of the product sold in this container being beyond the control of the seller, all conditions and warranties statutory or otherwise as to the quality or fitness for any purpose of such product are excluded and no responsibility will be accepted by the seller or manufacturer for any damage arising from it's application or use.

**ADDITIONAL SAFETY INFORMATION:**

Read label before use. May cause an allergic skin reaction. Wear protective gloves and clothing. If on skin wash with plenty of soap and water. If skin irritation or rash occurs get medical advice. Wash contaminated clothing before re-use. Toxic to aquatic life. Toxic to aquatic life with long lasting effects. Avoid release to the environment.

# SWAT

## INSECTICIDE FOR CATTLE

RAIN RESISTANT  
LOW-IRRITANT FORMULATION