



Product Name: Surefire Exclaim 44 SG Insecticide  
APVMA Approval No: 86238/115093

Label Name:	Surefire Exclaim 44 SG Insecticide
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	ACTIVE CONSTITUENT: 44 g/kg EMAMECTIN present as EMAMECTIN BENZOATE
Mode of Action:	GROUP 6 INSECTICIDE
Statement of Claims:	For the control of Diamondback Moth (Cabbage Moth) and Cabbage White Butterfly in Brassica Vegetables and Heliothis in Capsicum, Lettuce and Tomatoes
Net Contents:	NET CONTENTS: 500 g – 20 kg
Restraints:	
Directions for Use:	This section contains file attachment.
Other Limitations:	
Withholding Periods:	WITHHOLDING PERIODS Brassica Vegetables: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION DO NOT USE TREATED CROP, CROP WASTE OR PRODUCE FOR STOCK FOOD Capsicum, Lettuce, Tomatoes: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION DO NOT ALLOW LIVESTOCK TO GRAZE TREATED CROPS OR CROP STUBBLE

Trade Advice:	<p><b>EXPORT TRADE ADVICE</b></p> <p>Treated crop commodities destined for export may require extra time being allowed between application and harvest, as some export market have either no Maximum Residue Limits (MRLs) or different MRLs to those of Australia. Details of overseas standards and any export interval can be obtained by contacting PCT Holdings Pty Ltd before using this product.</p>
General Instructions:	<p><b>GENERAL INSTRUCTIONS</b></p> <p><b>MIXING</b> Add the required amount of Surefire Exclaim to a partly filled spray tank, and then add the remainder of the water.</p> <p><b>APPLICATION</b> Surefire Exclaim is not systemic making good coverage essential - ensure thorough coverage of the crop DO NOT apply by aerial application Brassica Vegetables: Apply Surefire Exclaim in a minimum of 400 L water/ha. Capsicums, Lettuce, Tomatoes: Ensure thorough coverage of the crop.</p> <p>Mixing/Application – Trellised Tomatoes Dilute Spraying: Use a sprayer designed to apply high volumes of water up to the point of runoff and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of runoff. Avoid excessive runoff. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of runoff. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed as the crop grows. Concentrate Spraying: Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of runoff) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (see Dilute Spraying) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way: Example Only 1. Dilute spray volume as determined above: For example 1000 L/ha 2. Your chosen concentrate spray volume: For example 500 L/ha 3. The concentration factor in this example is: <math>2 \times</math> (i.e. <math>1000 \text{ L} \div 500 \text{ L} = 2</math>) 4. If the dilute label rate is 15 g/100 L, then the concentrate rate becomes <math>2 \times 15</math>, that is 30 g/100 L of concentrate spray The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.</p>
Resistance Warning:	<p><b>INSECTICIDE RESISTANCE WARNING</b> <b>GROUP 6 INSECTICIDE</b></p> <p>For insecticide resistance management, Surefire Exclaim 44 SG Insecticide is a Group 6 Insecticide. Some naturally occurring insect biotypes resistant to Surefire Exclaim and other Group 6 insecticides may exist through normal genetic variability in any insect population. Repeated use of one particular insecticide or insecticides from the same chemical group can lead to the early development of insect resistance to that insecticide</p>

	<p>or chemical group. The effectiveness of Surefire Exclaim on resistant individuals could be significantly reduced. Resistant individuals can eventually dominate the insect population if Surefire Exclaim and other Group 6 insecticides are used repeatedly. Surefire Exclaim should not be applied more than two times per season and do not apply as consecutive applications. Since occurrence of resistant individuals is difficult to detect prior to use, PCT Holdings Pty Ltd accepts no liability for any losses that may result from the failure of Surefire Exclaim to control resistant insects. Surefire Exclaim may be subject to specific resistance management strategies. For further information contact your local supplier, PCT Holdings representative or local agricultural department agronomist.</p>
--	---

Precautions:	<p><b>PRECAUTION</b>  Re-entry Period: DO NOT allow entry into treated areas for 8 hours or until the spray has dried. If prior entry is necessary, wear chemical resistant gloves.</p>
--------------	---

Protections:	<p><b>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</b>  Dangerous to fish and other aquatic organisms. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT apply under weather conditions or from spraying equipment which could be expected to cause spray to drift onto adjacent areas, particularly wetlands, water bodies or watercourses.</p> <p><b>PROTECTION OF LIVESTOCK</b>  Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.</p>
--------------	---

Storage and Disposal:	<p><b>STORAGE AND DISPOSAL</b>  Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.  Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p>
-----------------------	---

Safety Directions:	<p><b>SAFETY DIRECTIONS</b>  Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing the product for use, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length PVC gloves and goggles or face shield. Wash hands after use. After each day's use, wash gloves, goggles or faceshield and contaminated clothing.</p>
--------------------	---

First Aid Instructions:	<p><b>FIRST AID</b>  If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.</p>
-------------------------	--

First Aid Warnings:	
---------------------	--

## DIRECTIONS FOR USE

CROP	PEST	RATE	CRITICAL COMMENTS
<b>Brassica Vegetables</b> Broccoli, Brussels Sprouts, Cabbages, Cauliflower only	Diamondback Moth (Cabbage Moth) ( <i>Plutella xylostella</i> ) and Cabbage White Butterfly ( <i>Pieris rapae</i> )	250 to 300 g/ha	Spray at first signs of insect infestation as indicated by local spray thresholds. Use the lower rate on low to moderate infestations. Add Citowett* or Agral® at the appropriate label rate. DO NOT make more than 4 applications to any brassica crop. Where more than one crop is grown DO NOT make more than 4 applications of Surefire Exclaim in any one year. Surefire Exclaim should be used according to CropLife insecticide resistance strategies.
<b>Capsicum, Lettuce</b>	Heliothis ( <i>Helicoverpa armigera</i> and <i>Helicoverpa punctigera</i> )	150 to 250 g/ha	Spray at first signs of insect infestation as indicated by local spray thresholds. For best results apply soon after the <i>Helicoverpa</i> eggs have hatched. DO NOT apply more than 4 sprays of Surefire Exclaim per crop. Where more than one crop is grown per year DO NOT apply more than 4 sprays per year. Ensure thorough spray coverage. Use the lower rate when targeting light infestations of small larvae. Use 250 g/ha during periods of heavy insect pressure or under very hot and sunny conditions. Surefire Exclaim should be used according to the CropLife resistance strategies.

CROP	PEST	RATE	CRITICAL COMMENTS
<b>Tomatoes</b>	Heliothis ( <i>Helicoverpa armigera</i> , <i>Helicoverpa punctigera</i> )	150 to 250 g/ha or 15 to 25 g/100 L (dilute spraying) for trellised Tomatoes. <b>Concentrate Spraying</b> Refer to the Mixing/ Application section	Spray at first signs of insect infestation as indicated by local spray thresholds. For best results apply soon after the <i>Helicoverpa</i> eggs have hatched. DO NOT apply more than 4 sprays of Surefire Exclaim per crop. Where more than one crop is grown per year DO NOT apply more than 4 sprays per year. Ensure thorough spray coverage. Use the lower rate when targeting light infestations of small larvae. Use 250 g/ha during periods of heavy insect pressure or under very hot and sunny conditions. <b>Trellised Tomatoes: In the case of dilute spraying (g/100 L) apply in the range of 400 to 500 L/ha after transplanting and increase to 800 to 1000 L/ha at full canopy.</b> Surefire Exclaim should be used according to the CropLife resistance strategies.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**